

Original Article

A Partial-Thickness Quadriceps Autograft Reliably Augments the Size of the Hamstring Graft During Anterior Cruciate Ligament Reconstruction

Yoan Bourgeault-Gagnon, M.D., Alexandre Keith Leang, M.D., F.R.C.S.C., Sonia Bédard, Karina Lebel, Ph.D., Frédéric Balg, M.D., F.R.C.S.C., and François Vézina, M.D., F.R.C.S.C.

Purpose: To measure the increase in diameter resulting from the augmentation of a hamstring autograft with a partial width rectus femoris tendon band in anterior cruciate ligament reconstruction. **Methods:** Thirty-three cadaveric knees were dissected to harvest semitendinosus and gracilis tendons (4S) along with a 6-mm wide tendon band from the rectus femoris. Harvesting was done according to the usual surgical techniques of both harvests. Measures of length and diameter in 4S and 4S augmented with the rectus femoris band (4S +Q) configurations were performed separately by 3 evaluators. **Results:** The quadriceps augmentation led to an average increase of 1.49 mm (95% confidence interval 1.03-1.95 mm) in diameter of the 4-strand hamstring grafts. The previously demonstrated threshold diameter of 8.5 mm was attained in only 30% of 4S grafts within this population in comparison with 88% when augmented with a quadriceps band. **Conclusions:** In conclusion, supplementing doubled hamstring graft (4S) with quadriceps tendon in anterior cruciate ligament reconstruction (ACLR) increases the graft diameter by an average of 1.49 mm. It has the physical potential to reliably augment hamstring grafts that measure 7.5 mm in diameter or more in order to obtain an 8.5 mm when necessitated. **Clinical Relevance:** Increased graft diameter is associated with a decreased risk of graft failure after ACLR. Because of this, it is important to identify methods to increase the size of grafts. This study investigates the use of a partial-width rectus femoris tendon band as an option to reliably augment graft sizes during ACLR.

Despite several advances in surgical techniques, the failure rate of anterior cruciate ligament (ACL) reconstruction generally varies between 5% and 25%.¹ Principal factors influencing these results are the definition of failure, the patient's age, the surgical techniques employed, the level of physical activity, and the size of grafts.¹⁻⁸

One of the most frequently used ACL reconstruction grafts is an autograft composed of doubled semitendinosus and gracilis (4-strand hamstring graft, or 4S).^{9,10} Biomechanical studies show an inversely proportional relationship between the graft's size and the risk of rerupture.¹¹ A large-scale clinical study relates a reduction in relative risk of almost 15% per 0.5-mm increase in graft diameter.^{12,13} A correlation between graft size and failure rate is demonstrated in multiple studies, with cutoffs established at 8 or 8.5 mm of diameter. These results are in agreement with those of other researchers who found a greater failure rate^{1,3,4,14} and lesser functional results⁴ with grafts measuring 8 mm in diameter or less.

Different options can be considered when the harvested graft's diameter is inferior to the target value. The most frequently used, due to its simplicity and the fact that it doesn't increase morbidity, is to triple the graft to obtain 5 or 6 strands (5S or 6S).^{15,16} However, in some situations, length could be unsatisfactory even after tripling the graft or the graft could, even tripled, maintain a smaller diameter than desired. In these cases, another supplementation technique might have to be added.

From Sherbrooke University (Y.B.-G., K.A.L., S.B., K.L., F.B., F.V.); Center for Research at the CHUS (CIUSSS de l'Estrie CHUS) (S.B., F.B.); and Research center on aging (CIUSSS de l'Estrie CHUS) (K.L.), Sherbrooke, Québec, Canada.

The authors report the following potential conflicts of interest or sources of funding: Y.B.-G., S.B., F.B., and F.V. report unrestricted grants from DePuy Synthes, A Johnson & Johnson Company, Wright Medical Company, and Zimmer Biomet, outside the submitted work. Full ICMJE author disclosure forms are available for this article online, as [supplementary material](#).

Received May 18, 2022; accepted August 24, 2022.

Address correspondence to Dr. François Vézina, Sherbrooke University, 3001 12th Avenue North, Sherbrooke, J1H 5H4, (Quebec), Canada. E-mail: francois.vezina@usherbrooke.ca

© 2022 THE AUTHORS. Published by Elsevier Inc. on behalf of the Arthroscopy Association of North America. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).
2666-061X/22543

<https://doi.org/10.1016/j.asmr.2022.08.009>

The option of a hybrid allograft/autograft is associated with a greater failure rate and a lesser integration than a 4S graft of the same diameter.¹⁷⁻¹⁹ In this context, one of the authors started using a partial width quadriceps tendon band (rectus femoris only) as a supplementation technique when a hamstring autograft has a diameter deemed unsatisfactory. Harvesting a quadriceps tendon band in ACL reconstruction is well-established and globally accepted as a principal graft.²⁰⁻²⁹ However, Wilson et al.³⁰ are currently the only authors having described and tested the biomechanical characteristics of using quadriceps tendons (Q) as supplementation of a 4S graft. The advantage of this technique relies on the ability to maintain the full length of the 4S graft. Only an additional 3-cm incision is needed to harvest this additional graft.

It is desirable to be able to estimate the impact of harvesting a band of quadriceps tendon on the graft's total diameter to ensure a sufficient size after supplementation. The purpose of this study was to measure the increase in diameter resulting from the augmentation of a hamstring autograft with a partial width rectus femoris tendon band in ACL reconstruction. We hypothesized that this technique would reliably increase the diameter of the graft by more than 1 mm.

Methods

The cadaveric specimens used in this study are part of a larger project. The data from this study constitute only a part of all the data collected from the specimens dedicated to the larger project. The study protocol was approved by the local ethical committee.

The specimens were prepared according to the usual technique for fresh corpse preservation, without embalming, by the technicians of the Medical Faculty's Anatomy Laboratory. A total of 34 cadaveric knees were dissected by one of the orthopaedic surgery senior residents to harvest the semitendinosus, gracilis, and quadriceps tendons while adhering to the usual surgical techniques. Of those 34, 33 samples were deemed usable in the study due to their quality, the absence of structural damage, and the absence of previous surgeries on those structures. The age and sex of the cadavers from which these samples were harvested have been tabulated (Table 1).

Harvesting of every tendon used was done by the same senior resident by using the instruments and surgical techniques typically employed in the authors' hospital. An oblique anteromedial incision is made near the insertion of the pes anserinus to the proximal tibia. Subcutaneous adipose tissues are dissected until the sartorius is seen, then incised longitudinally along the axis of its fibers. The gracilis tendon is then isolated and freed from its adherences before being harvested with the tendon harvester for cruciate ligament reconstruction (Conmed Linvatec, Aurora, OH). These steps are

Table 1. Demographic Characteristics

Number of samples	17 cadavers (33 samples)
Average age, y	76.2 ± 13.6
Body mass index	22.6 ± 4
Sex, men	52.9%

the same when harvesting the semitendinosus tendon. The remaining muscular fibers are cleaned from the tendons and the 2 tendons are freed in a subperiosteal manner from their tibial insertion, then set aside for ulterior measures. Harvesting of the quadriceps tendon band is done through a midline longitudinal incision reaching to the proximal pole of the patella. The subcutaneous adipose tissues are dissected until the quadriceps fascia is seen, then incised along the axis of its fibers with a 6-mm width in the central part of the tendon. The tendon band is then released from the patella with a depth of about 5 mm (thickness of the rectus femoris tendon) and this plane is dissected up proximally before being cut to a length of 9 cm. Each group of tendons from the same knee was identified with a code, then frozen until the day of the measurements.

At the time of measuring, each group of tendons had been unfrozen at room temperature and kept humid with surgical sponges soaked with normal saline between each step. The 4S configuration was then prepared for each group of tendons folded on a polydioxanone suture, and the diameter measures were taken using ACL diameter measuring tubes (Conmed Linvatec), varying from 6.5 to 11.0 mm in diameter in intervals of 0.5 mm. The graft length was measured with a millimeter graduated ruler. The quadriceps tendon bands were then sutured at the center of the 4S grafts to produce grafts in a 4S+Q conformation (Fig 1), which were also measured with the Conmed measuring tubes.

Each measure was done individually by 3 evaluators and compiled by a research assistant. Each evaluator




Fig 1. 4S+Q graft configuration. Four-strand hamstrings graft with quadriceps augmentation configuration. Black arrow-head: Quadriceps band. White arrow: Four-bundle hamstring graft.

Table 2. 4S and 4S+Q Graft Diameters

Diameter, mm	No. of 4S, %	No. of 4S+Q, %
7.0	9 (27.3%)	0 (0%)
7.5	6 (18.2%)	0 (0%)
8.0	8 (24.2%)	4 (12.1%)
8.5	9 (27.3%)	4 (12.1%)
9.0	1 (3.0%)	8 (24.2%)
9.5	0 (0%)	8 (24.2%)
10.0	0 (0%)	7 (21.2%)
10.5	0 (0%)	1 (3.0%)
11.0	0 (0%)	1 (3.0%)

4S, 4-strand hamstring graft; 4S+Q, 4-strand hamstring graft with quadriceps tendon supplementation.

also re-evaluated 10 grafts, randomly assigned by the research coordinator, to establish intraobserver reliability. It is worth noting that the repeated measures were done in a blinded manner. Further analysis of the impact of the augmentation technique were performed using the median value obtained from the three evaluators, for each sample.

Statistical calculations and analyses were done by a statistician with SPSS (version 23.0.0 from IBM Corp., Armonk, NY). Descriptive statistics were used for the calculation of means, medians, proportions, and confidence intervals. Inter- and intraobserver reliability of graft diameter measures was evaluated with the help of Kendall's coefficient of concordance, which allows evaluation of the concordance of discrete ordinal measures.

Results

The demographic characteristics of the studied cadavers can be found in Table 1. Intraobserver reliability, calculated for 4S+Q grafts, and interobserver reliability, calculated for all the measures in all configurations, are both excellent, with concordances of 0.976 ($P = .002$) and 0.959 ($P < .001$) respectively.

The diameter measures used in the results for each sample constitute the median of the three evaluators.

Table 3. Diameter Increase with 4S+Q Graft Augmentation

4S Diameter	Mean Increase (95% CI)	Percentage of Specimens of at Least 8.5 mm After Augmentation in 4S+Q	
7	1.50 (1.14-1.87)	66.67%	} 82,6% } 92,9%
7.5	1.33 (0.92-1.65)	83.33%	
8	1.38 (1.14-1.62)	100.00%	
8.5	1.56 (1.26-1.86)	100.00%	
9	1.0	100.00%	
Global	1.49 (1.03-1.95)	100.00%	

4S+Q, 4-strand hamstring graft with quadriceps tendon supplementation; CI, confidence interval.

Within this population, 69.7% of grafts had a diameter inferior to 8.5 mm with a 4S configuration (Table 2), whereas that ratio fell to 12.1% after supplementation with a 6-mm band of rectus femoris tendon (Table 2).

Supplementation with a quadriceps tendon band saw an average increase of 1.49 (95% confidence interval 1.03-1.95) in diameter for grafts in 4S conformation (Table 3). After augmentation, 82.6% of grafts with a diameter of 7 mm to 8 mm reached the target of 8.5 mm, in comparison with 92.9% of those with a diameter of 7.5 mm to 8 mm.

Discussion

The results of this descriptive study confirm a significant increase, of more than 1 mm, of the diameter of a 4S graft when supplemented with a quadriceps tendon band. Starting a few years ago, the senior author of this document used this 4S+Q supplementation technique as a backup option when the 4S graft diameter was insufficient, and its length did not allow to triple the hamstring tendons in a 6S configuration while using the same tibial fixation. Being part of a larger project aimed at creating a decisional algorithm for graft selection according to diameter and length, this study focuses on better establishing the boundaries and criteria for the use of this type of supplementation. It states that 92.86% of grafts with a 7.5 mm or more diameter that did not reach the target of 8.5 mm in diameter with a 4S conformation did so with a 4S+Q conformation with the advantage of maintaining the full length of the doubled hamstring graft.

We realize that adding a second harvesting site adds potential complications. Based on our experience, the 2 main potential complications following the harvest of a quadriceps tendon band—namely some pain and the postoperative strength of the extensor mechanism³¹—do not seem to be major stakes in the recuperation of patients benefiting from this supplementation. Existing literature demonstrates significantly less anterior knee pain^{23,28,32-34} and less kneeling pain^{23,28,32,35} after harvesting quadriceps tendons when compared with patellar tendon, even when it is harvested with the patellar bone and in its entire thickness. Although it remains to be demonstrated in a subsequent study, we believe that with a quadriceps tendon harvesting technique of partial thickness only, of reduced width and without any bone block, these complications are even rarer. Also, new minimally invasive harvesting techniques could further reduce these potential complications.³⁶

This descriptive study backs the use of quadriceps tendon supplementation described by Wilson et al.³⁰ and constitutes only the second study on the matter. In this sense, it contributes in establishing the foundations necessary to the elaboration of further clinical projects.

Limitations

The fact that this study is based on cadavers and that the mean age of the donors doesn't represent the usual ACL tear population can diminish the clinical applicability of its results. The absolute diameter of the tendons used can differ from the one in our aimed population because of age and preservation process. The total number of samples ($n = 33$) directly limits the numbers per group and might therefore influence the external validity of the experiment. Several questions, such as the complications associated with the technique as well as the real clinical benefit, have not been evaluated at all and are beyond the scope of this project.

Conclusions

In conclusion, supplementing doubled hamstring graft (4S) with quadriceps tendon in ACL reconstruction increases the graft diameter by an average of 1.49 mm. It has the physical potential to reliably augment hamstring grafts that measure 7.5 mm in diameter or more in order to obtain an 8.5 mm when necessitated.

Acknowledgments

Special thanks to Claudia Beaulieu, Sonia Paquette, and Denis Bisson, technicians in the anatomy laboratory, who helped us greatly with the preservation and preparation of samples. We would also like to thank Catherine Raynaud, who helped with data collection, and Sean David Gagnon, who helped for the English translation.

References

- Magnussen RA, Lawrence JTR, West RL, Toth AP, Taylor DC, Garrett WE. Graft size and patient age are predictors of early revision after anterior cruciate ligament reconstruction with hamstring autograft. *Arthroscopy* 2012;28:526-531.
- Shelbourne KD, Gray T, Haro M. Incidence of subsequent injury to either knee within 5 years after anterior cruciate ligament reconstruction with patellar tendon autograft. *Am J Sports Med* 2009;37:246-251.
- Conte EJ, Hyatt AE, Gatt CJ, Dhawan A. Systematic review hamstring autograft size can be predicted and is a potential risk factor for anterior cruciate ligament reconstruction failure. *Arthroscopy* 2014;30:882-890.
- Mariscalco MW, Flanigan DC, Mitchell J, et al. The influence of hamstring autograft size on patient-reported outcomes and risk of revision after anterior cruciate ligament reconstruction: A multicenter orthopaedic outcomes network (MOON) cohort study. *Arthroscopy* 2013;29:1948-1953.
- Borchers JR, Pedroza A, Kaeding C. Activity level and graft type as risk factors for anterior cruciate ligament graft failure: A case-control study. *Am J Sports Med* 2009;37:2362-2367.
- Salmon L, App B, Sci V, et al. Incidence and risk factors for graft rupture and contralateral rupture after anterior cruciate ligament reconstruction. *Arthroscopy* 2005;21:948-957.
- Mohtadi N, Chan D, Barber R, Paolucci EO. Reruptures, reinjuries, and revisions at a minimum 2-year follow-up: A randomized clinical trial comparing 3 graft types for ACL reconstruction. *Clin J Sport Med* 2016;26:96-107.
- Marchand JB, Ruiz N, Couprie A, Bowen M, Robert H. Do graft diameter or patient age influence the results of ACL reconstruction? *Knee Surg Sports Traumatol Arthrosc* 2016;24:2998-3004.
- Charalambous CP, Kwaees TA. Anatomical considerations in hamstring tendon harvesting for anterior cruciate ligament reconstruction. *Muscles Ligaments Tendons J* 2012;2:253-257.
- Hammer DL, Brown CH, Steiner ME, Hecker AT, Hayes WC. Hamstring tendon grafts for reconstruction of the anterior cruciate ligament: Biomechanical evaluation of the use of multiple strands and tensioning techniques. *J Bone Joint Surg Am* 1999;81:549-557.
- Boniello MR, Schwingler PM, Bonner JM, Robinson SP, Cotter A, Bonner KF. Impact of hamstring graft diameter on tendon strength: A biomechanical study. *Arthroscopy* 2015;31:1084-1090.
- Snaebjörnsson T, Hamrin Senorski E, Ayeni OR, et al. Graft diameter as a predictor for revision anterior cruciate ligament reconstruction and KOOS and EQ-5D values: A Cohort study from the Swedish National Knee Ligament Register based on 2240 patients. *Am J Sports Med* 2017;45:2092-2097.
- Spragg L, Chen J, Mirzayan R, Love R, Maletis G. The effect of autologous hamstring graft diameter on the likelihood for revision of anterior cruciate ligament reconstruction. *Am J Sports Med* 2016;44:1475-1481.
- Park SY, Oh H, Park S, Lee JH, Lee SH, Yoon KH. Factors predicting hamstring tendon autograft diameters and resulting failure rates after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthroscopy* 2013;21:1111-1118.
- Krishna L, Panjwani T, Ren Mok Y, Keng Lin Wong F, Singh A, Jie Toh S. Use of the 5-strand hamstring autograft technique in increasing graft size in anterior cruciate ligament reconstruction. *Arthroscopy* 2018;34:2633-2640.
- Brown CH. Editorial Commentary: How to increase hamstring tendon graft size for anterior cruciate ligament reconstruction. *Arthroscopy* 2018;34:2641-2646.
- Wang H-D, Gao S-J, Zhang Y-Z. Comparison of clinical outcomes after anterior cruciate ligament reconstruction using a hybrid graft versus a hamstring autograft. *Arthroscopy* 2018;34:1508-1516.
- Perkins CA, Busch MT, Christino M, Herzog MM, Willimon SC. Allograft augmentation of hamstring anterior cruciate ligament autografts is associated with increased graft failure in children and adolescents. *Am J Sports Med* 2019;47:1576-1582.
- Pennock AT, Ho B, Parvanta K, Edmonds EW, Chambers HG, Roocroft JH, et al. Does allograft augmentation of small-diameter hamstring autograft ACL

- grafts reduce the incidence of graft re-
tear? *Am J Sports Med* 2017;45:334-338.
20. Hughes JD, Vaswani R, Gibbs CM, Tisherman RT, Musahl V. Anterior cruciate ligament reconstruction with a partial-thickness quadriceps tendon graft secured with a continuous-loop fixation device. *Arthrosc Tech* 2020;9:e603-e609.
 21. Kanakamedala AC, de SA D, Obioha OA, et al. No difference between full thickness and partial thickness quadriceps tendon autografts in anterior cruciate ligament reconstruction: A systematic review. *Knee Surg Sports Traumatol Arthrosc* 2019;27:105-116.
 22. Stäubli HU, Schatzmann L, Brunner P, Rincón L, Nolte LP. Mechanical tensile properties of the quadriceps tendon and patellar ligament in young adults. *Am J Sports Med* 1999;27:27-34.
 23. Cavaignac E, Coulin B, Tscholl P, et al. Is quadriceps tendon autograft a better choice than hamstring autograft for anterior cruciate ligament reconstruction? A comparative study with a mean follow-up of 3.6 years. *Am J Sports Med* 2017;45:1326-1332.
 24. Lee S, Seong SC, Jo H, Park YK, Lee MC. Outcome of anterior cruciate ligament reconstruction using quadriceps tendon autograft. *Arthroscopy* 2004;20:795-802.
 25. Rabuck SJ, Musahl V, Fu FH, West RV. Anatomic anterior cruciate ligament reconstruction with quadriceps tendon autograft. *Clin Sports Med* 2013;32:155-164.
 26. DeAngelis JP, Fulkerson JP. Quadriceps tendon—a reliable alternative for reconstruction of the anterior cruciate ligament. *Clin Sports Med* 2007;26:587-596.
 27. Diermeier T, Tisherman R, Hughes J, et al. Quadriceps tendon anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2020;28:2644-2656.
 28. Slone HS, Romine SE, Premkumar A, Xerogeanes JW. Quadriceps tendon autograft for anterior cruciate ligament reconstruction: A comprehensive review of current literature and systematic review of clinical results. *Arthroscopy* 2015;31:541-554.
 29. Hurley ET, Calvo-Gurry M, Withers D, Farrington SK, Moran R, Moran CJ. Quadriceps tendon autograft in anterior cruciate ligament reconstruction: A systematic review. *Arthroscopy* 2018;34:1690-1698.
 30. Wilson WK, Morris R, Coskey A, Smith B, Gugala Z. Quadriceps augmentation of undersized hamstrings during ACL reconstruction. *Knee* 2019;26:73-78.
 31. Iriuchishima T, Ryu K, Okano T, Suruga M, Aizawa S, Fu FH. The evaluation of muscle recovery after anatomical single-bundle ACL reconstruction using a quadriceps autograft. *Knee Surg Sports Traumatol Arthrosc* 2017;25:1449-1453.
 32. Gorschewsky O, Klakow A, Pütz A, Mahn H, Neumann W. Clinical comparison of the autologous quadriceps tendon (BQT) and the autologous patella tendon (BPTB) for the reconstruction of the anterior cruciate ligament. *Knee Surg Sports Traumatol Arthrosc* 2007;15:1284-1292.
 33. Han HS, Seong SC, Lee S, Lee MC. Anterior cruciate ligament reconstruction: Quadriceps versus patellar autograft. *Clin Orthop Rel Res* 2008;466:198-204.
 34. Lund B, Nielsen T, Faunø P, Christiansen SE, Lind M. Is quadriceps tendon a better graft choice than patellar tendon? A prospective randomized study. *Arthroscopy* 2014;30:593-598.
 35. Kim SJ, Kumar P, Oh KS. Anterior cruciate ligament reconstruction: Autogenous quadriceps tendon-bone compared with bone—patellar tendon—bone grafts at 2-year follow-up. *Arthroscopy* 2009;25:137-144.
 36. Slone HS, Xerogeanes JW. Anterior cruciate ligament reconstruction with quadriceps tendon autograft: A minimally invasive harvest technique. *JBJs Essent Surg Tech* 2014;4:e16.